



Department
for Environment
Food & Rural Affairs

Biosecurity and preventing welfare impacts in poultry and captive birds

Advice for all captive bird and poultry keepers (including game birds, waterfowl, and pet birds)

14 December 2020



Llywodraeth Cymru
Welsh Government



Scottish
Government
gov.scot

Contents

Introduction	3
Benefits of biosecurity	3
Spread of disease	3
Biosecurity measures	4
Buying new stock	7
Other actions to consider for all poultry keepers	7
Protecting your birds from wild birds	8
Housing	9
Netted structures	10
Feeding and watering	11
Deterrents for wild birds	11
Pheasants, partridges and other game birds	11
Domestic ducks and geese	11
Captive wildfowl and waterfowl	12
Backyard flocks and pet/hobby birds	12
Zoological collections of birds	13
Planning permission for bird housing	14
Steps to take before letting birds outside (This only applies when the AIPZ is revoked) ...	14
Effects on organic and free-range status	15
Welfare: taking care of your birds	16
Welfare surveillance	16
Maintaining welfare after the release of free-range hens from housing (This only applies when the AIPZ is revoked)	18
Racing pigeons	19
Birds of prey	20
Further help and advice	21
What to do if you find dead wild birds	21
Be vigilant!	21
Advice and guidance	21

Introduction

This document tells you, as a keeper of poultry or captive birds, about the biosecurity measures you should take to prevent your birds from becoming sick with diseases like avian influenza and Newcastle disease. You should read it if you keep birds for commercial, hobby or other purposes, including as pets.

You should read this document together with the specific legal requirements in the Avian Influenza Prevention Zones that England, Scotland and Wales declared on 11 November 2020, and extended on 14 of December 2020 to include a housing requirement. The purpose of the zones is to help protect poultry and other captive birds within Great Britain from a number of highly pathogenic strains of avian influenza (bird flu) circulating in wild birds in the UK and Europe.

To assist you with this guidance, a [biosecurity self-assessment checklist](#) is provided. It is advised that you complete the self-assessment form, in order to assess and assure yourselves whether your current biosecurity practices are fit-for-purpose, and to identify where improvements may be needed. It has been designed to support you in meeting the measures introduced as part of an Avian Influenza Prevention Zone, in particular.

Keepers in England, Scotland and Wales **are required** to keep captive birds, chickens, turkeys and most other species of poultry housed indoors. For gamebirds and commercially farmed ducks and geese, where housing is less practicable due to welfare concerns, keepers must take steps to keep them separate from wild birds.

This document represents biosecurity guidance that is published in accordance with section 6A of the Animal Health Act 1981 in England and Wales. In Scotland, this document represents guidance that is published in accordance with the Codes of Recommendations for the Welfare of Livestock: Animal Health and Biosecurity, made under the Animal Health and Welfare (Scotland) Act 2006. You should read this document together with the specific declarations and legal measures in force at the time, and detailed on [GOV.UK](#), on [GOV.SCOT](#), or on [GOV.WALES](#). Measures could include Protection Zones, Surveillance Zones, Restricted Zones or an Avian Influenza Prevention Zone.

Benefits of biosecurity

Good biosecurity improves overall flock health and productivity by helping to keep out poultry diseases such as avian influenza and Newcastle disease. In the event of an outbreak, it also limits the spread of disease both on and off your premises and cuts the costs of disease treatment and reduces losses, which could improve profitability.

Spread of disease

Disease can be spread through several pathways, including:

- the introduction of diseased birds
- contact with diseased flocks
- through wild birds' contaminated faecal matter around the farm (such as walking it into sheds on footwear or on clothing and equipment)
- vermin or wild birds getting access to poorly maintained houses, feed or bedding
- birds drinking from contaminated water sources
- birds eating contaminated feed
- contamination of vehicles, equipment, clothing and footwear which are moved between and within farms
- using shared farm equipment and vehicles, which have not been effectively cleansed and disinfected
- unsatisfactory cleansing and disinfection of vehicles, sheds, feeding troughs or other equipment

Biosecurity measures

The following are mandatory biosecurity measures required within an Avian Influenza Prevention Zone to reduce the risk of transmission of highly pathogenic avian influenza. All poultry keepers (including game birds, waterfowl, and pet birds) must take all appropriate and practicable steps to ensure that:

- **You wear clean overalls and footwear when entering poultry farms or premises, to avoid bringing infection onto your farm, or spreading it around farms or premises, via your clothes, footwear or hands. Overalls should ideally be changed and foot baths used when moving between flocks or sheds.**
- **Staff and all visitors should be provided with protective clothing and footwear dedicated for use on your premises.** These should be removed and cleansed, disinfected, laundered or disposed of after use. Remember to include maintenance men or workmen and any tools they bring onto the premises. Provide hand-washing facilities, and insist that these are used by staff and all visitors.
- **Keep good records of all visitors and vehicles that enter the site** – names, contact details, dates, times and purpose of visit. This will allow the source and spread of disease to your premises to be rapidly traced and will minimise the effect on your business.
- **Strictly limit and control access to all poultry flocks (including game birds and pet birds). If possible, the site should be fenced with a controlled entry**

point. You should restrict the number of visitors and their vehicles, and keep them as far away as possible from poultry buildings, and pastures.

- **Keep appropriate records of poultry (including game birds and pet birds), captive birds and egg movements including production records.** You must keep mortality records. You are also advised to record details of any movement of poultry and eggs on and off your premises, and relevant production records at a shed/house level (such as water consumption, feed consumption and egg production). Where possible, records should be electronic.
- **Have pressure washers, brushes, hoses, water, and fresh supplies of an [approved disinfectant](#) available at all points where people should use them.** Make sure they are used by all visitors to clean vehicles, equipment and boots both before entry to, and upon leaving, your farm and poultry houses or enclosures. Visitors should, ideally, only use footwear and outer clothing that is kept on-site.
- **Make sure that disinfectant in boot dips is at the right concentration and that it is changed in accordance with the manufacturer's instructions.** Use a stable disinfectant with good resistance to inactivation by organic matter, at a minimum concentration as required by the disinfectants approved for use in England, Scotland and Wales (<http://disinfectants.defra.gov.uk>). The level of boot dip should always be at least ankle deep and care should be taken to position the boot dip to allow easy step in/step out procedures to take place. Boot dips should be sited in sheltered/covered locations to prevent disinfectant from being diluted by rainwater. The effectiveness of the disinfectant can become compromised in the presence of excessive organic material. Therefore, they should be checked regularly and if this occurs, the disinfectant should be changed. Pre-cleaning boots with water and a brush prior to using the dip is essential to avoid organic overload. It is preferable to have specific footwear that is only worn in the bird area, changing footwear on entering and leaving the area accessed by birds.
- **Cleanse and disinfect all vehicles which have been transporting poultry (including game birds and pet birds), poultry products or poultry by-products, after each journey.** Don't forget to clean the wheels, under wheel arches wheels, and foot wells.
- **Cleanse and disinfect all crates, containers, plastic egg trays and other equipment before and after use or if not possible, destroy.** Do not move any equipment into different poultry buildings without cleansing and disinfecting it first. This also applies to injecting and dosing equipment.
- **Keep farm access routes, parking areas, yards, areas around buildings and storage areas clean and tidy and well maintained, at all times.** This helps avoid wild birds and animals being attracted onto your premises and entering buildings and stores. It will also reduce the risk of vehicles becoming

contaminated. You should also prevent the accumulation of standing water and remove any spilled feed that could attract wild birds.

- **Keep all poultry houses or sheds well maintained to ensure that wild birds do not nest or roost in them, and cannot access the inside of the building.** Rectify any defects, such as holes in the roof or walls. Any openings must be covered with suitable mesh or netting which should be of a maximum size of 25 mm mesh to prevent ingress from wild birds (See the advice on protecting your birds from wild birds.)
- **Keep all poultry houses or sheds well maintained to ensure that water cannot ingress.** Rectify any defects, such as holes in the roof or walls or damaged guttering, without undue delay to prevent ingress of contaminated water.
- **Keep wild birds, dogs, cats, rodents and other livestock out of poultry buildings and feed stores.** These can carry infection or infective material onto your farm and spread disease to your flock.
- **Have an effective rodent and pest control system in place.** Be vigilant for evidence of vermin. Monitor vermin activity by baiting and trapping. Feed silos and containers must be regularly cleaned, maintained and properly sealed, to prevent vermin and wild birds accessing and contaminating feed.
- **Supply only clean fresh drinking water to birds.** Water lines and drinkers must be flushed through and cleaned regularly. You could consider the use of water sanitisers in consultation with your vet. Where birds cannot be housed for welfare reasons restrict access to possible sources of water used by wild birds and restrict wild bird access to your drinking systems.
- **You should only obtain your feed from a mill or supplier that operates in accordance with relevant Government, Agricultural Industries Confederation or credible assurance scheme standards,** who will normally make available results of salmonella tests on request.
- **Damaged eggs, dead birds, litter and manure may carry disease. Dispose of them promptly and appropriately.** For further information, see the guidance on fallen stock and on poultry manure:
 - [Fallen stock and safe disposal of dead animals in England](#)
 - [Fallen stock guidance in Scotland](#)
 - [Fallen stock guidance in Wales](#)
 - [How to use, store or move manure, guano and digestive tract content](#)

- **At depopulation at the end of a production cycle (for example at the end of lay for laying hens), thoroughly clean the building and all equipment, including ducting, drains and fans.** Remove all surplus feed, dead birds and litter. Disinfect the premises and all equipment and carry out rodent and other pest control. You should also clean and disinfect cleaning equipment and protective clothing. Make sure that any visiting teams that catch your poultry follow the biosecurity measures described above for entering and leaving your premises, and only wear protective clothing that is dedicated for use on your premises. If partial depopulation (for example, on a broiler unit - thinning) is carried out, make sure a quarantine period is observed by catching teams. A quarantine of 72 hours is recommended between their last contact with poultry and entry to your buildings.

Buying new stock

- Check whether poultry and/or captive bird gatherings are permitted. If they are not, then buying new stock from a gathering will not be possible (although direct sales that are not at a gathering may still go ahead).
- Always make sure you know the health status of any birds you are buying or moving.
- Incoming stock should be isolated from the rest of the flock – discuss this with your vet and agree a testing and monitoring programme.
- Only place new stock in facilities which you know have been cleansed and disinfected.
- Where used, keep isolation buildings as near as possible to the farm entrance and separate from other poultry buildings/ranges.
- Use separate equipment when handling isolated stock. If you have enough staff on your farm, allocate some of them to deal only with isolated stock. If not, make sure you handle isolated stock last. Always wash and change into clean overalls and boots before going back to your main flock buildings.

Other actions to consider for all poultry keepers

Where practical, you should also take the following steps:

- Consult your private veterinary surgeon about the risks specific to your premises and the practical steps you can take to reduce them.

- For poultry flocks with fewer than 50 birds, although you are not required by law to register them, we encourage you to do so, as this means we can contact you quickly if there is an outbreak of disease. Please see further information on [how to register your poultry](#).
- Keep domestic ducks and geese completely separate from other poultry species as they often don't show any signs of disease but can still pass it on to chickens, turkeys or other poultry species.
- If you have staff, give them the information and training they need to maintain good standards of hygiene (an important aspect of biosecurity) at all times. If they keep their own birds at home, they will need to be extra vigilant for signs of disease and even more careful about biosecurity both at home and on your farm.
- You must consider what measures may be appropriate and practical to protect your birds from contact with wild birds, such as feeding and watering birds indoors or under protection from wild birds and rodents (see the advice on protecting your birds from wild birds later in this guidance). Talk to your private veterinarian for further advice.
- Review the contingency plans which you should have in place for managing your premises in the event of a suspected or confirmed case of notifiable avian disease, both on your premises or nearby. The types of movement restrictions that could be put in place are set out in the [Notifiable Avian Disease Control Strategy](#). Register your premises with APHA on the GB Poultry Register. You should also regularly check that APHA have your correct contact details, including email address and mobile phone number. This will allow us to contact you in the event of an outbreak in your area or further afield in Great Britain. It will help in avoiding delays in issuing licences in the event of an outbreak.
- Sign up to the [Animal and Plant Health Agency \(APHA\) Diseases Alerts Service](#) to keep up to date with the latest news on other exotic notifiable animal disease outbreaks in Great Britain. Note, this is not the same as the GB Poultry Register.

Protecting your birds from wild birds

One way by which notifiable avian disease may spread to poultry is through contact with infected wild birds. Contact may be direct (through mingling), or indirect (through bird secretions, faeces, feathers, rodents etc., contaminating anything that may then come into contact with poultry such as feed, water, utensils clothing or vermin). Flooding at your premises or fields nearby can attract wild birds, thereby increasing the risk of notifiable avian disease. Flood water can be contaminated (with wild bird infectious material), which can get into poultry houses or range areas.

Make sure poultry houses or sheds on your premises are well maintained to prevent

wild birds from nesting or roosting in them. It is important that you regularly inspect the building(s) used to house your poultry (including game birds and pet birds) for any structural damage (such as leaks e.g. roofs, holes, blocked drains or downpipes) and repair any defects without undue delay. Notifiable avian disease can be introduced into poorly maintained poultry houses, via wild birds, rats or mice that could mechanically carry virus into the house, or ingress of contaminated water. It is your responsibility to make sure your birds are protected from disease causing agents.

Biosecurity is cumulative, so the more you do, the better protected your birds will be.

Where an Avian Influenza Prevention Zone or other disease control zone such as a Protection Zone, Surveillance Zone or Restricted Zone has been declared, the declaration will state the mandatory biosecurity and wild bird separation measures that will apply. **You must follow the disease control measures and any published guidance in force at the time.**

Housing

Due to an increased risk of incursion of Avian Influenza to domestic poultry and wild birds within you **must** take all appropriate and practicable steps to ensure that your birds are housed, or otherwise kept separate from wild birds. Chickens and turkeys are especially susceptible to Avian influenza and it is expected that all keepers will house their birds. Housing your birds is the most effective way to minimise direct contact with wild birds, provided that the housing does not have entry points that can be exploited by wild birds. Housing of birds on its own is not a substitute for the best practice biosecurity measures described above.

You must make sure that wild birds are prevented from gaining access to your birds by **covering any gaps, openings or uncovered ranges with netting**. This should have a mesh size that is sufficiently small enough to exclude wild birds from passing through the openings (see the section below for advice on mesh size). Prior to 14 December 2020, if your birds spent time outside you must consider alternative arrangements. Planning permission may be required for some of these options and the section on planning (below) should be consulted. Check these structures regularly for damage that may allow wild birds access or ingress of water. There are a number of different housing options, including:

- making use of suitable existing buildings (such as barns, farm sheds, outbuildings, garages, garden sheds) adapted for your birds. Ensure that there is adequate ventilation and that any openings or ranges are netted to prevent wild birds gaining access. You should check for and remove hazardous and toxic substances if garages and outbuildings are being used as temporary accommodation.

- erecting a solid lean-to or veranda on the side of existing houses (provided it does not impact on the minimum space requirement for access to outdoor areas for free range birds).
- erecting a new temporary structure with solid walls and roof, or a polytunnel. A polytunnel will be suitable only in cooler weather.
- any proposed building must take into account the welfare requirements of the birds, in particular the importance of adequate ventilation and light. This is also covered in the advice on welfare later in this guidance.

Netted structures

Total netting or full enclosure is required as an alternative to mandatory housing for birds that cannot easily be housed; for example, ducks and geese may be kept in fully netted areas, even if this means netting a smaller area than they currently occupy. (The only exception is for certain zoo/aquarium birds and overwintering birds kept for restocking game, where other measures to minimise contact with wild birds should be used).

Netting of fenced areas can also reduce the risk of disease, providing wild birds cannot gain access to the enclosed area and steps are taken to make the area unattractive to wild birds.

If you choose to fully enclose/totally net outdoor areas, the following steps should be taken:

- erecting a net structure that will reduce contact between your birds and wild birds;
- when using a netted structure, pay special attention to keeping wild birds from perching on the roof and defecating through it - for example you should consider using additional deterrent methods described below
- constructing temporary outdoor pens using straw bales and a tarpaulin roof with bird-proof netted gaps for light and ventilation
- if bad weather is likely to be a problem, it may be necessary to erect a windbreak around your bird-proof structure
- it is generally recommended that netting should have a maximum mesh size of 25mm although for exposed areas or where heavy snow fall can be expected, a larger mesh size of up to 50mm may be appropriate. You should, where appropriate, consult commercial suppliers of anti-bird netting for advice and further information on equipment

Feeding and watering

There are certain species of bird – such as ostrich, over-wintered poultry kept for the restocking of game, which are not normally housed, for which the housing steps outlined above may not be practicable. In such cases, you must isolate their food and water from wild birds. Available feed and water will attract wild birds; by feeding and watering your birds under cover, and ensuring feed spills and leaks from silo augers etc. are prevented, the possibility of mingling is reduced. The steps you should take, where practical, include:

- never feed or water your birds outdoors in open troughs or by scattering feed on the ground. Feed and water should be provided under cover or in specially designed feeding stations that prevent access by wildbirds.
- rotating feeding times. Many wild birds learn when kept birds are fed and congregate at these times.
- preventing your birds from accessing open water that may be contaminated. Ensure that your birds receive only mains or treated water, or ensure that reservoirs or storage tanks are covered. Sealed nipple systems can be considered.

Deterrents for wild birds

When designing any protective structure, if you need to use posts, you should think about incorporating spike strips to deter perching by wild birds. In all cases you should also consider making use of deterrents such as flutter tape, flashing lights and scarecrows.

However, in doing so, you should be careful to avoid any potential impacts on your birds; particularly flashing lights.

Pheasants, partridges and other game birds

Birds should be housed whenever possible, or kept in fully netted areas where practical. All feeding and watering should take place under cover. Specialist advice is available from representative organisations and the Animal and Plant Health Agency (APHA). For overwintering birds kept for the restocking of game, where housing or netting is not possible, other methods of preventing contact with wild birds, such as providing feed and water under cover and other methods of deterring wild birds, should be used.

Domestic ducks and geese

Where small numbers of ducks and geese are kept, it may be possible to house them.

However, if that is not possible, ducks and geese should be kept in fully netted areas or temporary netted structures, even if it means confining them to a smaller area. All feeding and watering should take place indoors or under cover. You should consider using wild bird deterrents to prevent contact with wild birds. Ducks and geese should not be kept with other poultry species as they often don't show any signs of disease, but can still pass it on to chickens, turkeys or other poultry species.

Captive wildfowl and waterfowl

Birds should be housed whenever possible or otherwise kept separate from wild birds in fully netted areas as outlined by an AI Prevention Zone. All feeding and watering should take place under cover or in some form of structure designed to exclude wild birds as far as possible. Consider making sensible use of wild bird deterrents to prevent contact with wild birds. You should consider moving your birds away from large bodies of water that attract wildfowl.

Where separation is not possible, for example where there are large numbers of waterfowl, every effort should be made to discourage wild birds and keep feed separate.

Backyard flocks and pet/hobby birds

All keepers, regardless of how many birds they keep must meet these requirements. Birds **must** be housed in a suitable building, such as a specialist poultry house, shed, outbuilding or garage adapted to house birds, or a new temporary structure such as a lean-to or polytunnel (note - a polytunnel will be suitable only in cooler weather). Ensure openings and ranges are netted to prevent wild birds getting in and remove any hazardous substances. Anyone who keeps chickens, turkeys, quail and fancy fowl is expected to house their birds, and those with other poultry (including but not limited to guinea fowl, geese and ducks) should fully enclose or net their outdoor runs and pens.

You are responsible for ensuring the welfare of your birds while they are indoors, and for keeping them calm and comfortable:

- ensure birds have natural light where possible and are not kept permanently in the dark. Any artificial light should ideally follow typical day and night patterns
- ensure there is adequate ventilation – adult birds will tolerate low temperatures but are very susceptible to high temperatures (temperatures should not be allowed to go significantly above 21°C)
- feather pecking can be a risk when birds are kept indoors. Keep the environment interesting to reduce this risk. Consider adding pecking blocks, fresh bedding, straw bales and objects such as cabbages, scatter feed or whole grain on the

floor of the building and add grit to litter so birds can scratch and carry out natural foraging behaviors

- if feather pecking occurs, you can reduce light levels, although they should still be light enough for you to read a newspaper
- if you keep several types of birds, you should house chickens or turkeys in separate enclosures from waterfowl (ducks and geese)
- check birds regularly to ensure they are healthy and have enough food, water and dry bedding; you may want to consider nutritional supplements in drinking water as these can help keep birds calm
- skin parasites, such as red mite, can be a problem in birds kept indoors and can make birds more irritable. Advice on controlling these parasites can be obtained from your vet
- [Register your birds on the GB Poultry Register.](#)

Zoological collections of birds

It is recognised that licensed zoos and aquariums house a wide range of bird species, often of high conservation value, and that it is not always possible to ensure the basic welfare requirements of particular species can be met by housing or full netting.

Therefore, subject to agreement with APHA and the British and Irish Association of Zoos and Aquariums (BIAZA), zoos (a premises licensed under the Zoo Licensing Act 1981.) with captive birds do not need to fully net outside enclosures.

Unless there is a justifiable welfare concern, zoos must adhere to the biosecurity requirements set out in the AIPZ declaration. Deviation from these requirements will only be allowed in exceptional circumstances. The zoo, in consultation with their vet, will be expected to produce robust justification for any deviation and demonstrate they have measures in place to minimise any disease spread from these un-netted outdoor enclosures.

Zoos may choose to manage their operations as a single unit or split their site into multiple units. The unit should contain the area in which the birds are kept and keeper areas (changing area, area to keep PPE and equipment used in the enclosure, cleaning and disinfection facilities, storage of waste pending appropriate disposal). Each unit can then be treated as if it were a separate premise and the biosecurity measures outlined in an AIPZ Declaration applied to each in turn.

If the site is split into multiple units, zoos must demonstrate that these are epidemiologically distinct, that is, there is no transfer of birds, food, equipment or personnel between the different units, without them having undergone suitable

disinfection, decontamination / isolation / quarantine etc. Zoos should consult their veterinary surgeon about how they can split their site into epidemiological units and what measures they need to put in place to keep these units separate.

Zoos also have to recognise the biosecurity risks introduced from access by members of the public and steps should be taken to minimise the risk of disease transmission to and from zoological collections by visitors. Where members of the public are permitted access to areas where poultry or other captive birds are kept, disinfectant mats should be placed at entry and exit points, and handwashing facilities made available.

All zoos should have a written, site specific, disease contingency plan produced in consultation with their vet.

Planning permission for bird housing

Planning permission for bird housing is not always required if the work does not involve building, engineering or similar operations. Permitted Development Rights exist for erecting structures in certain circumstances. Simple structures such as posts and netting might not require permission. It is unlikely, for example, that you would need planning permission if the structure is temporary, mobile, or used for the purposes of agriculture.

However, regulations will differ between commercial and hobby poultry keepers. You should always check with your local planning authority in the first instance. You can go to www.planningportal.gov.uk for contact details in England and Wales or [planning authorities in Scotland](#).

Steps to take before letting birds outside (This only applies when the AIPZ is revoked)

Before releasing birds after a period of housing, there are a number of steps you should take to prepare the outside area (the “range”) and reduce the risk of infection. To be effective these steps should be taken as early as possible before birds are allowed back onto the range:

- inspect your range regularly and remove any obvious contaminants from wild birds (such as faeces or feathers) in a biosecure manner
- net ponds and drain waterlogged areas of land. If this isn't possible, then you can fence them off from your birds so they cannot access it whilst ranging, or use an alternative paddock that doesn't have access to water
- remove any feeders and water stations from the range, or ensure that they are

covered to sufficiently restrict access by wildbirds

- consider using decoy predators on the range, or allowing dogs to accompany you on foot patrols around the range. You could also consider bird scarers if their use is appropriate for the area (see [NFU Code of Practice](#) on bird scarers)
- consider daily checks of the range area to monitor any changes in the biosecurity measures in place and to record changes in wild bird activity and discuss with your own vet where these are noted
- consider increasing the number of shelters on the range area

Effects on organic and free-range status

The legal requirement for keepers to house their birds or otherwise kept separate from wild birds as part of an Avian Influenza Prevention Zone, also applies to free-range and organic poultry producers. Chickens and turkeys are especially susceptible to Avian influenza and it is expected that all keepers will house their birds.

- The organic status of poultry flocks is not affected by any legal requirement to house or restrict access to open-air runs, provided that all other requirements of the [Organic Standards](#) continue to be met.
- From the 14th December 2020, eggs from poultry currently reared as free range can continue to be marketed as free range provided continuous housing does not exceed 16 weeks (in this case, up till, 00.01hours on 5th April 2021), if they meet all other requirements. The 16-week period applies to the time when eggs are laid, rather than the time of sale.
- From the 14th December 2020, poultrymeat from poultry currently reared as free range can continue to be marketed as free range provided continuous housing does not exceed 12 weeks (in this case, up till, 00.01hours on 8th March 2021), if they meet all other requirements. The 12-week period applies on a per-flock basis.
- For longer-term concerns about possible loss of status, for example organic or free-range, you should consult the responsible sector body or organic certification body.

Welfare: taking care of your birds

Welfare surveillance

- You must always look after the welfare of your birds, particularly so in the case of a disease outbreak. If you choose to house your birds for biosecurity or commercial reasons, or housing is required by government and your birds are not used to being housed for a sustained period, this may create welfare and behavioural problems. This means:
- you should seek advice from your vet if there are any potential concerns about welfare.
- you should be vigilant for signs such as injurious feather pecking, behaviours associated with redirected foraging and any other signs of poor welfare such as loss of condition
- you should understand the type of pecking that could occur in your flock. Aggressive pecking, which is not the same as injurious feather pecking, may also be seen when management changes are made. [Featherwel](#), BEIC and AssureWel have produced practical guidance which describes how to check for early signs of injurious feather pecking in laying hens. [AssureWel](#) has also produced a bench- marking tool for feather loss scores for commercial producers.
- you should consider using scratch feeds to prevent boredom amongst housed flocks, which can lead to injurious feather pecking and cannibalism
- overcrowding may become a problem and you must take steps to ensure that the stocking density within each house will not reach the point at which bird welfare is compromised – a particular risk in hot, cold or wet weather
- you should consider installing thermometers at bird level – you should monitor the maximum temperature, and take action if it significantly exceeds 21°C for adult birds. This could prevent losses due to hyperthermia (high temperatures are more difficult for birds to cope with than low temperatures), but they may also give an indirect measure of ventilation rates
- you should consider providing perches as a welfare benefit. In Scotland, for species other than waterfowl, it is a requirement that all Scottish registered laying flocks are provided with aerial perching. There are many advantages, including making use of vertical space so that stocking density is reduced, but the main advantage is giving submissive birds the chance to remove themselves from situations where they may otherwise be pecked or cannibalised
- you should provide fresh litter in all temporary accommodation, and ensure that adequate feed and water, ventilation and lighting are available. Consider making use of damp proof membranes for earthen floors to maintain good

litter conditions

- you should also consider ways to enrich the birds' environment more generally, such as using straw bales or hanging cabbages
- you should maintain a regular lighting pattern with no sudden changes. If an outbreak of injurious feather pecking occurs, you may need to reduce the lighting, as well as adding more environmental enrichment including scratch feeding if feasible
- always ensure that your birds have clean water – do not use standing water that may have become contaminated by wild birds
- you should also ensure that you (or your staff) inspect the birds more often in order to pick up problems before they lead to the deterioration of bird welfare
- you should avoid sudden changes to main diet and, where possible, mix old and new diets through any transition period. Mashing feed increases eating time compared to pellets and keeps the birds occupied which reduces the risk of injurious feather pecking. Consumption of whole oats, wheat, corn, alfalfa, maize/barley/pea, silage and carrots can reduce injurious feather pecking, plumage damage and death
- you should maintain effective levels of disease control. High levels of red mite infection and worm burdens are associated with increased risk of injurious feather pecking

you should provide pullets with rearing conditions that are similar to those during laying. If pullets are normally given access to a range during their rearing phase and are then confined, this can increase the risk of injurious feather pecking. Pullet suppliers should make any unexpected confinement history available to those taking delivery of these birds to help reduce the risk of injurious feather pecking at onset of lay and during the laying period

Specific measures for ducks

You may wish to provide an alternative source of water for housed ducks. Open water sources such as troughs, filled water buckets and showers enable them to immerse their heads in or under water and preen effectively, keeping eyes, nostrils, beaks and plumage in a healthy condition. However, your ducks should not be given access to water which may be contaminated by wild birds. You should ensure that litter quality does not deteriorate with excess water spillage. Wet bedding can increase the spread and severity of pododermatitis (contact dermatitis on the feet) and an increased susceptibility to other infectious diseases.

Consider making use of raised, perforated plastic floors or equivalents to place water sources on to assist maintaining litter/straw quality.

You should ensure that ventilation and temperature controls are appropriate for ducks. They drink and produce more water in their droppings than other poultry, and humidity and ammonia levels can increase rapidly. Adult ducks prefer lower temperatures (around 13°C) to other poultry.

Having a contingency plan that can be put into operation will make sure the welfare of your flock remains a priority at a critical time. Further guidance and information on poultry welfare is available on the following pages:

- England: [poultry on farm welfare](#), or you can call the Defra Helpline on **03459 33 55 77**
- Scotland: [farmed animals](#)
- Wales: [animal welfare](#)

Maintaining welfare after the release of free-range hens from housing (This only applies when the AIPZ is revoked)

When released after a period of housing, free-range poultry may exhibit increased levels of aggression or feather pecking. The cause of these behavioural changes is not completely understood, but should be treated in a similar manner to any occurrence of these behaviours. If you are worried about their welfare, seek advice from your vet.

You should continue to inspect livestock more regularly than once a day. This will ensure that you detect any injurious pecking resulting from housing at the earliest opportunity, and enable you to take steps to treat or cull injured birds. Ideally, hens should be kept inside large naturally-lit veranda-type buildings. If this is not possible, you should maintain light intensity at 10 lux or above in the perching, feeding and walking areas during the housed period and afterwards unless injurious pecking occurs. You should only decrease light intensities for the period where injurious pecking is a welfare problem.

During housing and afterwards, resources such as food and water sources, should be made more widely available to reduce competition.

Space allowances for free-range hens are specified in legislation and codes of recommendations and should be maintained at these levels or more.

Re-introduction to the range should be carried out with minimal disturbance to the

birds. You should treat the flock as if it was a young flock and re-introduce birds to the range gradually over a period of 7-10 days in order for them to re-adjust to temperature changes, light conditions and additional external influences.

Racing pigeons

Bird gatherings are not allowed to take place anywhere in Great Britain. You can fly pigeons that already live together, in an Avian Influenza Prevention Zone (AIPZ). However, birds that are not already housed together/on the same premises should not be flown together, as this constitutes a bird gathering.

You can exercise your racing pigeons for up to one hour once each day while an AIPZ with housing measures in force, but they should avoid direct contact with wild birds. You may train birds as a single loft, but must not mix birds from other lofts. You cannot do this if you are in a Protection Zone or Surveillance Zone.

You must not have an open loft that allows the birds to freely come and go, which would increase the chance of wild birds entering your loft and contaminating it.

To reduce the risk of your birds becoming infected with Avian Influenza, you should take steps to avoid your birds coming in to contact with wild birds, or their faeces/body fluids:

- You should focus on ensuring a high level of cleanliness (biosecurity) in your lofts, particularly if you also keep poultry – chickens, hens, ducks, geese, turkeys, game birds and guinea fowl – and should follow the advice in biosecurity guidance documents available on government websites.
- You should undertake vermin (rats and mice) control to reduce their contact with feed and water for pigeons.
- Your birds should, where practicable, receive mains or treated water that is not accessible to wild birds or poultry to reduce the risk of contamination.
- You should ensure your baskets are as clean as practicable when you are using them to transport birds to other locations (other lofts, shows, sales, race marking). They should be disinfected regularly during the racing season.
- You should consider cleaning the feet of your birds before transporting them to other locations, particularly for shows, races and sales.
- If you also keep other poultry, including a few chickens or ducks, ensure that your pigeons neither mix with such poultry nor have access to their food and water, in order to reduce the risk of spreading disease.
- Any vehicles / trailers used to transport pigeons must be cleansed and disinfected using Government approved disinfectant, between transports

Birds of prey

Bird gatherings are not allowed to take place anywhere in Great Britain. You can fly a bird (falcon/hawk etc.) or multiple birds that already live together, in an AIPZ with housing measures in force.

You can fly birds of prey, including for pest control, if they are already housed together, but they should avoid direct contact with wild birds. Birds that are not already housed together/on the same premises should not be flown together, as this constitutes a bird gathering. You cannot do this if you are in a Protection Zone or Surveillance Zone.

Further help and advice

What to do if you find dead wild birds

Do not touch or pick up any dead or visibly sick birds that you find in the wild. If you find dead wild waterfowl (swans, geese or ducks) or gulls, or any other dead wild birds of other species in the same location in England, Scotland or Wales, you should report them to the Defra helpline on **03459 33 55 77** or by emailing defra.helpline@defra.gov.uk. We may then collect some of these birds and test them to help us understand if disease is present and, if so, how this may be distributed geographically and in which bird species.

Where dead birds are not required for surveillance purposes it is the landowner's responsibility to safely dispose of the carcasses as animal by-products.

Be vigilant!

Look out for clinical signs of disease in your flock.

Increased mortality, unusual behaviour, neurological signs such as shaking or incoordination, decreased eating or drinking, falling egg production, eggs with thin or misshapen shells and breathing difficulties may all be early signs of a disease problem. Ensure you keep yourself and your staff aware of what signs each disease may cause.

If you suspect disease, ask your vet for advice as soon as possible. Do not wait for more evidence – **some diseases can spread very quickly.**

Where avian influenza (or Newcastle disease) is **not** strongly suspected, but cannot be ruled out, poultry keepers may wish to liaise with their private veterinarian about utilising the Animal and Plant Health Agency (APHA) "testing to exclude" regime in GB. This involves submitting samples to a testing service at the APHA's National Reference Laboratory, Weybridge. This can help detect a notifiable avian disease at the earliest opportunity, for such cases. Further information is available on the [APHA website](#).

Avian influenza and Newcastle disease are notifiable diseases and if you suspect either of these diseases, you must report them immediately by calling the Defra Rural Services Helpline on 03000 200 301. In Wales, contact 0300 303 8268. In Scotland, contact your local [Field Services Office](#). Failure to do so is an offence.

Advice and guidance

Always practice good biosecurity, ask your vet for advice if you suspect disease. You must report any suspicion of avian influenza or Newcastle

disease immediately by calling the Defra Rural Services Helpline on 03000 200 301. In Wales, contact 0300 303 8268. In Scotland, contact your local [Field Services Office](#). Failure to do so is an offence.

You can find more information about diseases, such as avian influenza and Newcastle disease, including signs to watch out for, at the following pages:

- England: [avian influenza](#) and [Newcastle disease: how to spot and report it](#), or you can call the Defra Helpline on **03459 33 5577**.
- Scotland: [avian influenza](#) and [Newcastle disease](#).
- Wales: [avian flu](#) and [Newcastle disease](#)

Version Control	Date
Version 1.0	Published 21 December 2014
Version 1.1	<p>Published on 29 November 2016.</p> <p>Minor editorial changes to biosecurity advice and updates to links. Additional information regarding the Animal and Plant Health Agency “testing to exclude” scheme included on page 12 and new paragraphs added on welfare surveillance on page 13.</p>
Version 1.2	<p>Published on 7 December 2016.</p> <p>Document updated to refer to specific biosecurity considerations applicable across Great Britain under Avian Influenza Prevention Zones that were created by Defra, the Scottish Government and the Welsh Government on 6 December 2016.</p>
Version 1.3	<p>Published on 18 January 2017</p> <p>Document updated to reflect the extension to the Prevention Zones to 28 February 2017, also includes extra guidance for those with backyard flocks.</p>
Version 1.4	<p>Published on 31 March 2017</p> <p>Document updated to remove specific references to any current Prevention Zones which may be in force, and to include extra guidance on welfare.</p>
Version 1.5	<p>Published on 29 May 2017</p> <p>Minor editorial changes, plus amendments to Housing subsections to make it clear that housing of birds is a voluntary decision, unless required by government.</p>
Version 1.6	<p>Published on 27 November 2018</p> <p>Minor editorial changes and updates to include keeping ducks and geese separate from other poultry.</p>
Version 1.7	<p>Published on 1 April 2020</p> <p>Minor editorial changes including amendments to general biosecurity subsections and “Protecting your birds” paragraph.</p>
Version 1.8	<p>Published on 11 November 2020.</p> <p>Document updated to refer to specific biosecurity considerations applicable across Great Britain under Avian Influenza Prevention Zones that were created by Defra, the Scottish Government and the Welsh Government on 11 November 2020.</p>

Version 1.9	Published on 27 November 2020 Minor editorial change to reflect the quarantine period needed to be observed by catching teams during partial depopulation.
Version 2.0	Published on 3 December 2020 Document updated to reflect the mandatory housing measures due to come into force on 14 December 2020. across Great Britain.
Version 2.1	Published on 14 December 2020 Document updated to reflect the mandatory housing measures coming into force on 14 December 2020 across Great Britain.



You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v.3. To view this licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/ or email PSI@nationalarchives.gov.uk

This publication is available at www.gov.uk/government/publications

Any enquiries regarding this publication should be sent to us at

Animal Health Policy, Exotic Diseases Control
Defra
Area 2D, Nobel House, 17 Smith Square
London SW1P 3JR

PB 11380